Case Report

Samarathunga AAS¹, Fernando C², Fernando J³

Introduction

Desmoplastic Infantile Astrocytoma (DIA) of the central nervous system is an uncommon intracranial neoplasm in the paediatric age group. It comprises solid areas of neoplastic astrocytes within a prominent desmoplastic stroma and large cystic spaces. This tumour characteristically involves the superficial cerebral cortex and leptomeninges in infants. It has been described by various names in early literature and is probably regarded as a form of malignant mesenchymal tumour (1,2). Taratuto and co-workers introduced the term meningo-cerebral astrocytoma attached to the dura, with desmoplastic reaction in 1982 (1). Vanden Berg et al described it as desmoplastic supratentorial neuroepithelial tumour of infancy with divergent differentiation in 1987 (2). Currently this tumor is recognized as a distinct entity and is included in the World Health Organization (WHO) classification under the category of neural and mixed glioneuronal tumors, in grade I of the WHO classification (3,4).

Case Report

A 5-month-old female was admitted with uncontrollable seizures and delayed milestones. She was the first child in the family born from normal vaginal delivery after an uncomplicated full term pregnancy.

Physical examination revealed increased head circumference for the age with no other neurological manifestations. Magnetic resonance imaging (MRI) scan showed a large hypodense area in the left parietal region with extensive white matter oedema, mild mid line shift and left side uncal herniation. A cystic and solid mass attached to the dura was excised via a left frontoparietal craniotomy. The post operative period was uneventful, with no seizures after surgery.

The specimen consisted of multiple fragments of grayish and firm tissue, altogether measuring about 40X20X20 mm. Microscopically, the haematoxylin and eosin (H&E) stained sections revealed a markedly desmoplastic tumor with dense collagen fibers. The neoplastic cell population was heterogeneous, composed predominantly of spindle shaped astrocytes with a fascicular arrangement. Scattered ganglion cells were also observed, indicating neuronal differentiation (Figure 1). No mitoses or necrosis was present. The tumour revealed a rich reticulin network. A preliminary diagnosis of benign/low grade desmoplastic spindle cell tumour was made. Immunoperoxidase stains were performed GFAP stain highlighted the neoplastic astrocytes within an unstained desmoplastic stroma while vimentin was diffusely positive (Figure 2).

^{1,3}*Department of Pathology, National Hospital of Sri Lanka, Sri Lanka.*

²*Department of Pathology, Faculty of Medicine, University of Colombo, Sri Lanka.*

Correspondence: Fernando J, Department of Pathology, National Hospital of Sri Lanka, Sri Lanka. e-mail: janae@sltnet.lk

Focal isolated ganglion cells were positive with synaptophysin and Neuron Specific Enolase (NSE), but EMA was negative.

Fig 1. Desmoplasia and fascicular pattern of the arrangement (H&EX10)

Fig 2. GFAP positive astrocytes and unstained desmoplastic component (GFAPX40)

Discussion

Desmoplastic infantile astrocytoma is a rare intracranial tumour that typically occurs in infants below the age of 24 months (3,4). Studies indicate an average incidence of about 1% of all a desmoplastic infantile astrocytoma (WHO grade I). Follow up of the patient after 2 months was unremarkable. paediatric tumours (4,5). These tumors are large in size and predominantly cystic, with preferential frontal and

Desmoplastic infantile astrocytoma; an uncommon paediatric intracranial tumour

parietal involvement, as seen in this patient. Boys are affected more commonly than girls (6). In contrast, this tumour was in a girl. Typically, a DIA appears as a hypointense cystic mass with an isointense peripheral solid component on MR images similar to this case. However, radiologic and histologic features of DIA are misleading, and it may be misinterpreted as a malignant lesion (5). In this patient MRI was reported as suspicious for a primitive neuroectodermal tumour (PNET). Symptoms of DIA are intracranial hypertension, sunset eyes, enlarging head circumference, bulging fontanelle, variable localizing signs including seizures or paresis (7). This patient presented with intractable seizures and delayed milestones.

Microscopic features of the tumour have been clearly described in several studies (1,2,3). The most prominent feature of DIA is the presence of desmoplasia and spindle cells with a storiform pattern of arrangement (5). There are also ganglion cells which is present as single cells or in clusters (2,5). The first component can be shown to be GFAP positive, but the latter component is of neuroepithelial origin and reactive with markers such as synaptophysin (8). This tumor revealed typical histological and IHC features of DIA.

The main histological differential diagnoses are reticulin rich desmoplastic tumors such as pleomorphic xanthoastrocytoma, which can be differentiated by age of the patient, prominent lipidization of the cells and absence of neural component (7). Another differential diagnosis is a ganglioglioma. However, the age of the patient and lack of desmoplasia in ganglioglioma will

differentiate it from DIA (7). Complete resection is usually curative with no further additional therapy (8).

References

1. Taratuto AL, Monges J, Lylyk P, Leiguarda R. Superficial cerebral astrocytoma attached to dura. Report of six cases in infants. *Cancer* 1984 December 1; 54(11): 2505-12.
2. Vanden Berg SR, May EE, Rubinstein LJ et al. Desmoplastic supratentorial neuroepithelial tumors of infancy with divergent differentiation potential ("desmoplastic infantile gangliogliomas"). Report on 11 cases of a distinctive embryonal tumor with favorable prognosis. *Journal of Neurosurgery* 1987 January; 66(1): 58-71.
3. Pommepuy, MDelage-Corre, JJ Moreau and F Labrousse. Clinical Study A Report of a Desmoplastic Ganglioglioma in a 12-year-old Girl with Review of the Literature. *Journal of Neuro Oncology* 2006; 76: 271-275.
4. Brat DJ, Vandenberg SR, Figarella-Branger D, Taratuto AL: Desmoplastic infantile astrocytoma and ganglioglioma. In: World Health Organization classification of tumors of the central nervous system, Louis DN, Ohgaki H, Wiestler OD, Cavenee WK, 4th ed. Lyon:IARC press 2007; 96-98.
5. Trehan G, Bruge H, Vinchon M et al. Imaging in the diagnosis of desmoplastic infantile tumor: retrospective study of six cases. *American Journal of Neuroradiology* 2004 June-July; 25(6): 1028-33.
6. Per H, Kontaş O, Kumandaş S, Kurtsoy A. A report of a desmoplastic non-infantile ganglioglioma in a 6-year-old boy with review of the literature. *Neurosurgery Review* 2009; 32: 369-74.
7. Khaddage A, Chambonniere ML, Morrison AL, Allard D, Dumollard JM, Pasquier B et al. Desmoplastic infantile ganglioglioma: A rare tumor with an unusual presentation. *Annals of Diagnostic Pathology* 2004; 8: 280-3.
8. Craver RD, Nadell J, Nelson JS. Desmoplastic infantile ganglioglioma. *Pediatric and Development Pathology* 1999; 2: 582-5.
9. VandenBerg SR. Desmoplastic infantile ganglioglioma and desmoplastic cerebral astrocytoma of infancy. *Brain Pathology* 1993; 3: 275-81.